



Evaluation of Major Adverse Cardiac Events after Coronary Artery Bypass Graft Surgery and Valve Surgery in Gaza

Mohammed Habib, MD, PhD and Abdelaziz abu Samara, MD

Cardiology Department, Alshifa Hospital, Gaza, Palestine



Abstract

Background: The present study assessed the major adverse cardiac events (MACE) (hospital mortality rate, major stroke and reintervention) after cardiac surgery due to coronary artery bypass grafting or valve surgery or both in the Ministry of health hospitals (MOH) in Gaza Strip at 2022 by either Local or Expert cardiac surgery team.

Method: Retrospective cohort study including all patients who underwent cardiac surgery by local or expert cardiac surgery team with cardiopulmonary bypass and/or valve surgery in the MOH hospitals at Gaza Strip during 2022.

Result: Among the 98 patients (27 female and 72 male) with mean age 56 ± 12 years, 38 patients was operated by local team, and 60 patients was operated by expert cardiac surgeon during 2022 in Gaza. The comparison of end points (mortality, major stroke and reintervention) between local and expert cardiac surgeons, was significantly low in patients operated by expert surgeon was 9 patients (23.7%) in local team surgery and in 4 patients (6.7%) in expert team (P 0:015).

Conclusions: The combination end points included Hospital mortality, major stroke and reintervention rate was higher in patients operated by local cardiac surgery team.

Introduction

Cardiovascular disease (CVD) is the leading cause of death in the world, accounting for approximately 17.5 million deaths worldwide every year, of which 80% occur in low- and middle-income countries. Coronary artery bypass grafting (CABG) has remained the most commonly performed cardiac surgery procedure for coronary artery disease over the past decades [1]. The main goal of the surgery is to improve survival and health related quality of life (QoL) [2]. Despite the fact that the profile of patients undergoing isolated CABG surgery has changed towards higher age with increased frequency of preoperative comorbidities [3,4] mortality rate and morbidity after CABG have even decreased over past decades in the world [5].

The purpose of the present study was to compare hospital mortality rates of patients seen in the MOH health system who underwent coronary bypass surgery or valve surgery for the first time in publicly hospitals by local or foreign cardiac surgeon at MOH hospitals during 2022.

Methods

Data collection

From 1st January 2022 to 31st December 2022, data were prospectively collected for all patients covered by the public

healthcare system in whom a coronary artery bypass graft (CABG) and/or cardiac valve surgery was indicated as first surgery. This was done at two publicly managed hospitals (Alshifa and European Gaza Hospital).

Type of surgery

First time coronary artery bypass graft (CABG) and/or cardiac valve surgery.

End Points (MACE)

Hospital mortality

It was defined as death due to any cause during or after the procedure and up to hospital discharge, or within 30 days following the procedure.

***Corresponding author:** Mohammed Habib, MD, PhD, Cardiology Department, Alshifa Hospital, Gaza, Palestine

Accepted: August 03, 2023

Published online: August 05, 2023

Citation: Habib M, Samara A (2023) Evaluation of Major Adverse Cardiac Events after Coronary Artery Bypass Graft Surgery and Valve Surgery in Gaza. J Cardiothorac Surg Ther 7(1):152-154

Reintervention

The event was defined as the first reintervention for acute coronary syndrome after cardiac surgery. This reintervention could be a cardio logical intervention such as a PCI or similar procedure on a cardio surgical intervention such as isolated or combined CABG, mitral valve repair or replacement for ischemic valve disease within 90 days.

Major stroke

Patients with permanent Disability and NIHSS (National Institute of Health Stroke Scale) NIHSS more than 15 and Modified Rank in Scale mRS > 2 within 30 days.

Expert cardiac surgeon

Working total operation at least > 1000 cases and emperce > 10 years after graduated, we have 3 expert cardiac surgeon 2 from USA and one from Germany who make operation as short period in MOH in Gaza.

Statistical analysis

Baseline variables Continuous data are reported as means \pm SD. Categorical data are presented as absolute values and percentages. Using the x2 test for calculation of MACE (mortality rate, major stroke and reintervention) between patients underwent cardiac surgery local and expert cardiac surgeon. Significance level was set at P value < 0.05. Statistical analysis was performed with SPSS Statistics, Version 23.0.

Results

Among the 98 patients (27 female and 72 male) with mean age 56 ± 12 years scheduled for surgery during the study period, 38 patients was operated by local team, and 60 patients was operated by expert cardiac surgeon during 2022 in Gaza MOH hospitals (Table 1).

The comparison of end points (mortality, major stroke and reintervention) between local and expert cardiac surgeons was significantly low in patients operated by expert surgeon (Table 2).

The major causes or reintervention (incomplete revascularization in three patients, severe stenosis at anastomosis area in one patient, wrong anastomosis in one patient).

Discussion

The Gaza strip need 350-400 case for coronary artery bypass grafting and 200-250 for cardiac valve surgery yearly [6].

Because the restricted siege of Gaza from 2006 until now. Cardiac surgery was started in Gaza at 2010 by visitor cardiac surgeon from USA and Germany for making and educated local team, but the 30 days mortality in patients who operated by local cardiac surgeon still high 8% (three patients from 38 patients) and reintervention within 3 months is high 13% because of incomplete revascularization ang graft failure. Despite local team only operated selected low risk cardiac patients with elective ischemic coronary artery disease another cardiac surgery such as (valve surgery,

Table 1: Characterizes of patients.

Total patients	98
Mean age	56 + 12 years
Male	72 (73.5%)
Risk factors	
Hypertension	62 (63%)
Diabetes mellitus	43 (43.9%)
Type of Surgery	
CABG	74 (74%)
Valve surgery	22 (22%)
Combination (CABG + Valve)	2 (2%)
End Points	
Death	6 (6%)
Reintervention	5 (5%)
Major stroke	2 (2%)

Table 2: The comparison of MACE.

	Local (38)	Expert (60)	P Value
End points	9	4	0;015
Mortality	3	3	
Major strike	1	1	
Re-intervention	5	0	

aortic aneurysm, cardiac tumor, Redo CABG, Congenital heart disease surgery and CABG with acute coronary syndrome) still transfer to cardiac surgery hospitals in west bank.

In Canada, There is a trend to decreasing mortality in the AVR + CABG cohort to less than 3.0%, with isolated CABG and isolated AVR mortality relatively constant at 1.3% each [7].

In Gaza only two cardiac surgeon are available for 2.5 million people. A total of 12,180 adult cardiac surgeons were listed in the CTSNet registry, which translates to 1 adult cardiac surgeons per 0.61 million people globally, or, conversely, 1.64 adult cardiac surgeons per million people. Regional distribution ranged from 11.12 adult cardiac surgeons per million population (32.82% of total) in North America to 0.12 adult cardiac surgeons per million (1.05% of total) in sub-Saharan Africa [8].

Conclusion

Only two adult cardiac surgeon in Gaza is available, with limited surgery for coronary artery bypass grafting in patients with low risk and high mortality and reintervention rate after surgery. Gaza need additionally at least 2 to3 expert adult cardiac surgeon to make large number of different cardiac surgery and to reduce post-cardiac surgery complications.

References

1. D Agostino RS, Jacobs JP, Badhwar V, et al. (2019) The society of thoracic surgeons adult cardiac surgery database: 2019 update on outcomes and quality. The Annals of Thoracic Surgery 107: 24-32.

2. Neumann FJ, Sousa-Uva M, Ahlsson A, et al. (2019) 2018 ESC/EACTS guidelines on myocardial revascularization. *European Heart Journal* 40: 87-165.
3. Ferguson TB Jr, Hammill BG, Peterson ED, et al. (2002) A decade of change--risk profiles and outcomes for isolated coronary artery bypass grafting procedures, 1990-1999: A report from the STS national database committee and the duke clinical research institute. *The Annals of Thoracic Surgery* 73: 480-489.
4. Cornwell LD, Omer S, Rosengart T, et al. (2015) Changes over time in risk profiles of patients who undergo coronary artery bypass graft surgery: The veterans affairs surgical quality improvement program (VASQIP). *JAMA Surg* 150: 308-315.
5. ElBardissi AW, Aranki SF, Sheng S, et al. (2012) Trends in isolated coronary artery bypass grafting: An analysis of the society of thoracic surgeons adult cardiac surgery database. *J Thorac Cardiovasc Surg* 143: 273-281.
6. Habib Md, Adwan Md, Radi Md, et al. (2019) Incidence and trends of ischemic heart disease and coronary revascularization procedures in Gaza: A descriptive retrospective study. *Saudi Journal of Medicine* 4: 361-365.
7. Canadian Cardiovascular Society (CCS)/Canadian Society of Cardiac Surgeons (CSCS) Cardiac Surgery Quality Working Group. Discharge Abstract Database, National Ambulatory Care Reporting System and Hospital Morbidity Database, 2013-2014, 2014-2015, 2015-2016, 2016-2017 and 2017-2018, Canadian Institute for Health Information.
8. Vervoort D, Meuris B, Meyns B, et al. (2020) Global cardiac surgery: Access to cardiac surgical care around the world. *The Journal of Thoracic and Cardiovascular Surgery* 159: 987-996.e6.

DOI: 10.36959/582/448