



Research Article

DOI: 10.36959/468/487

Does CA-125 Have a Role in Early Diagnosis of Ovarian Malignancy in Non-Menopausal Women?

Momen Z El Nadeim¹, Bahgat KY Ahmed¹, Hanan H Mowad², Nesreen AA Shehata^{1*}  and Sara A Salem¹

¹Obstetrics and Gynecology Department, Faculty of Medicine, Beni-Suef University, Egypt

²Clinical and chemical pathology Department, Faculty of Medicine, Beni-Suef University, Egypt



Abstract

Objective: Aim of this study is to assess the role of cancer antigen (CA-125) in detection of ovarian malignancy in premenopausal women with ovarian mass.

Methods: This observational study was carried out in (blinded). It included 200 women who had preliminary diagnosis of adnexal mass whatever its nature. Adnexal masses were detected either clinically or by ultrasound.

Results: The study reveals of CA-125 cutoff point 35.1 and sensitivity and specificity reached 93.1% and 92.2% respectively. Area under the curve (AUC) was 0.998, Positive prediction value (PPV) was 91.7% and prediction value negative (NPV) was 92.1 (Table 1).

Conclusion: According to our results CA -125 could be suitable as an ovarian cancer detection marker.

Keywords

CA -125, Ovarian mass, Ovarian malignancy, Premenopausal women

Introduction

Ovarian cancer is one of the three most common malignant tumors in the female reproductive system. It has an insidious onset with a difficult early diagnosis [1]. In approximately, 70% of all cases of ovarian cancer, the disease is not diagnosed before reaching an advanced stage [2]. The 5-year survival rate associated with ovarian cancer is < 30% [3]. Over 90% of cases of ovarian masses detected in premenopausal and ≤ 60% in postmenopausal women are benign [4]. The early diagnosis of ovarian malignant tumor becomes a key factor in improving the survival rate of patients. Tools currently in use for differentiating between low-and high-risk patients with ovarian cancer are the tumor markers like cancer antigen-125 (CA-125) [5].

The tumor marker CA-125 has been used for 30 years for the monitoring of ovarian cancer, diagnosis, effective evaluation, and recurrence [6]. Although clinical application of CA-125 has been extensive, its specificity as a marker of malignant tumor or early diagnosis of ovarian cancer requires reassessment [7]. In premenopausal women, the detection of CA-125 in ovarian cancer sensitivity and specificity is not ideal because of the menstrual cycle, pregnancy and other effects [8]. Moreover, as there are no definite screening tools for ovarian malignancy and many pros and cons of tumor

markers regarding their sensitivity and specificity? We specify our search in this study on the CA-125 level and its role in ovarian cancer detection due to its high sensitivity, non-invasiveness, and simplicity [3,6,7].

Thus, we are assessing in this study the role of CA-125 in detection of ovarian malignancy in premenopausal women with ovarian mass.

Patients and Methods

An observational Cohort study was carried out in (blinded). The study was conducted from 2016-2018, according to the guidelines for good clinical practice for research and

***Corresponding author:** Nesreen Abd El Fattah Abd Allah Shehata, Obstetrics and Gynecology Department, Faculty of Medicine, Beni-Suef University, Egypt, Tel: 00-2010-2415-0605

Accepted: April 19, 2022

Published online: April 21, 2022

Citation: El nadeim MZ, Ahmed BKY, Mowad HH, et al. (2022) Does CA-125 Have a Role in Early Diagnosis of Ovarian Malignancy in Non-Menopausal Women?. Annals Gynecol Obstet 6(1):167-170

Table 1: Roc curve analysis of CA-125.

Area	Cutoff	P	Sensitivity	Specificity	PVP	PVN	95% Confidence Interval	
							Lower Bound	Upper Bound
0.998	35.1	0.00**	93.1%	92.2%	91.7	92.1	0.986	1.000

Table 2: Demographic data in between the study group.

Variable	
Age: (Years):	
Mean ± SD	37.76±11.68
Age groups	N (%)
25-39	100(50)
40-45	17 (8.5)
46-49	83 (41.5)
Parity :	
Median	2
Range	(0-4)

Table 3: Distribution of cases according to incidence of malignancy.

Incidence of malignancy	No. (%)
Malignant	54 (27)
Benign	146 (73)

declaration of Helsinki. Premenopausal women with adnexal masses participated in the study. All participants signed an informed consent form submitted for approval by the Ethical Review Board of the faculty of medicine, (blinded). The study included (200) premenopausal women who had preliminary diagnosis of an adnexal mass which was detected clinically and by ultrasound scanning.

Women were recruited from the outpatient gynecological clinic. After signing an informed consent, all participants were subjected to the following:

1. Full history taking with special focus on patient's age, parity, present history of the adnexal mass, family or past history of adnexal masses.

2. Blood sample for CA-125: Serum CA-125 level was determined by radioimmunoassay (MINIVEDAS CA-125 machine).

VIDAS® CA 125 II™ (125) VIDAS CA 125 II is an automated quantitative test for use on the VIDAS family instruments, for the measurement of OC 125 antigenic determinants in human serum or plasma (lithium heparin or EDTA) using the ELFA technique (Enzyme Linked Fluorescent Assay).

Sample size calculation: assuming that premenopausal women with ovarian mass attending (blinded) University Hospital was 280 and positive predictive value of CA-125 was 80.1, so the total sample was 200 women, using Epi-info at power 80% and CI 95%.

Results

The mean age of the studied cases was 37.76 ± 11.68 years. Median parity was 2 with a range of (0-4) (Table 2).

The study shows that 27% of cases had malignant tumors and while benign tumors were diagnosed in 73% of cases (Table 3).

The study reveals of CA-125 cutoff point 35.1 and sensitivity and specificity reached 93.1% and 92.2% respectively. Area under the curve (AUC) was 0.998, Positive prediction value (PPV) was 91.7% and prediction value negative (NPV) was 92.1 (Table 1).

Discussion

Currently, CA-125 is frequently used to detect ovarian cancer before the onset of clinical signs, but CA-125 can increase in association with some physiological conditions such as pre-menopausal women and benign diseases in women suspicious of cancer. There are other negative points about CA-125 biomarker properties which are, its low sensitivity for early-stage detection, and low specificity related to ovarian cancer. High level of CA-125 in the other cancers such as endometrial, cervix, and lung cancers is reported [9].

In our study the mean age of the studied cases was 37.76 ± 11.68 years. Median parity was 2 with a range of (0-4). These results are in agreement with study by Moore et al who they reported that the mean age for premenopausal women was 39.7 years [10].

Malignant epithelial ovarian tumors account for 90% of all malignancies of the ovary and are the fourth most common cause of tumor-related death in women [11].

In our study, 27% of cases had malignant tumors while 73% of cases had benign tumors.

Van Gorp, et al. in 2011 investigated 389 women: 228 (58.6%) patients had benign disease and 161 (41.4%) patients had malignant disease [12].

According to Partheen, et al. in 2011, their study population (n=374) included women with benign ovarian tumors (n=215), borderline type tumors (n=45), and epithelial ovarian cancer (EOC; n=114) [13].

In current study the cutoff point of CA-125 is 35.1 and sensitivity and specificity reached 93.1% and 92.2% respectively. Our results are supported by findings reported in a meta-analysis by Ferraro, et al. in 2013. They found that the specificity of CA125 for detecting ovarian cancer was 78% (95% CI 76-80) [14]. To describe tumor markers and screening tests, the Receiver Operating Characteristic (ROC) and area under the curve (AUC) are frequently employed since they represent a useful graphic tool for comparing biomarkers and algorithms. The ROC measures the discrimination of a test, i.e. its ability to distinguish between having disease and not having it for a given patient. In the study by Dikmen, et al. in 2015, the AUC for CA125 was rather weak (0.78), suggesting that it was probably not the ideal marker for diagnosing ovarian cancer [15].

Moore, et al. in 2010 included borderline tumors in their analysis, within this study, the examination of benign cases versus all stages of epithelial ovarian cancer and borderline tumors revealed a ROC-AUC of 0.913. Within a setting of a multicenter prospective trial with central review and monitoring it seems plausible that a diagnostic test would perform slightly better [16]. CA125 is higher in healthy premenopausal patients [17]. These slightly higher normal values influence the performance of the tumor markers concerned. Although not significant, this can also be seen in a study by Van Gorp, et al. in 2011, the ROC-AUC of CA125 was higher in the post-menopausal group [14].

Our results are supported by a multicenter clinical trial validating the performance of HE4, CA125 that suggesting that CA125 is superior to HE4 as a biomarker to detect ovarian cancer [18].

Anton, et al. in 2012 reported that the sensitivity value for CA125 detection was 83.8% with a specificity of 71.1%, whereas these values were 70.4% and 74.2%, respectively, when the tumors were classified as high-risk [15].

In 2011 from an analysis of patients with ovarian cancer, Chang X, et al. evaluated 491 patients and obtained a sensitivity of 88% using the marker CA125 [19].

In contrast, according to Oranratanaphan, et al. in 2018, HE4 and ROMA compared to CA-125, had lower sensitivity and NPV, but higher specificity and PPV for differentiating between benign and malignant ovarian tumor [20]. This result was consistent with that of the previous studies by Molina, et al. in 2011 and Chan, et al. in 2013 that were performed in 6 Asian countries including Thailand [21,22].

Furthermore, Roy in 2019 reported that sensitivity of CA 125 in the pre-menopausal women was 88.23% and that of the post-menopausal women was 100%. Specificity of CA 125 in the pre-menopausal women was 75.55% and that of the postmenopausal women was 88.88%. The positive predictive value in the pre-menopausal women was 57.69% and that of the post-menopausal women was 90%. The negative predictive value in the pre-menopausal women was 94.44% and that of the post-menopausal women was 100% [23].

In conclusion, application of the CA-125 measurement for the diagnosis of ovarian cancer was found to be effective and it has good clinical application, which is useful for clinicians.

References

1. Smith LH, Oi RH (1984) Detection of malignant ovarian neoplasms: A review of the literature. I. Detection of the patient at risk; clinical, radiological and cytological detection. *Obstet Gynecol Surv* 39: 313-328.
2. Heintz AP, Odicino F, Maisonneuve P, et al. (2006) Carcinoma of the fallopian tube. FIGO 26th Annual Report on the Results of Treatment in Gynecological Cancer. *Int J Gynaecol Obstet* 95: S145-S160.
3. Lee H, Kim JW, Lee DS, et al. (2020) Combined poziotinib with mandipine treatment suppresses ovarian cancer stem-cell proliferation and stemness. *Int J Mol Sci* 21: 7379.
4. Enakpene CA, Omigbodun AO, Goecke TW, et al. (2009) Preoperative evaluation and triage of women with suspicious adnexal masses using risk of malignancy index. *J Obstet Gynaecol Res* 35: 131-138.
5. Karlsen MA, Sandhu N, Høgdall C, et al. (2012) Evaluation of HE4, CA125, risk of ovarian malignancy algorithm (ROMA) and risk of malignancy index (RMI) as diagnostic tools of epithelial ovarian cancer in patients with a pelvic mass. *Gynecol Oncol* 127: 379-383.
6. Urban N, McIntosh MW, Andersen M, et al. (2003) Ovarian cancer screening. *Hematol Oncol Clin North Am* 17: 989-1005.
7. Nolen B, Velikokhatnaya L, Marrangoni A, et al. (2010) Serum biomarker panels for the discrimination of benign from malignant cases in patients with an adnexal mass. *Gynecol Oncol* 117: 440-445.
8. Folk JJ, Botsford M, Musa AG (1995) Monitoring cancer antigen 125 levels in induction chemotherapy for epithelial ovarian carcinoma and predicting outcome of second-look procedure. *Gynecol Oncol* 57: 178-182.
9. Bashizadeh-Fakhar H, Rezaie-Tavirani M, Zali H, et al. (2018) The diagnostic value of serum cea, ca-125, and roma index in low-grade serous ovarian cancer. *International Journal of Cancer Management* 11.
10. Moore RG, Miller MC, Disilvestro P, et al. (2011) Evaluation of the diagnostic accuracy of the risk of ovarian malignancy algorithm in women with a pelvic mass. *Obstet Gynecol* 118: 280-288.
11. Liest AL, Omran AS, Mikiver R, et al. (2019) RMI and ROMA are equally effective in discriminating between benign and malignant gynecological tumors: A prospective population-based study. *Acta obstet gynecol Scand* 98: 24-33.
12. Van Gorp T, Cadron I, Despierre E, et al. (2011) HE4 and CA125 as a diagnostic test in ovarian cancer: Prospective validation of the risk of ovarian malignancy algorithm. *Br J Cancer* 104: 863-870.
13. Partheen K, Kristjansdottir B, Sundfeldt K (2011) Evaluation of ovarian cancer biomarkers HE4 and CA-125 in women presenting with a suspicious cystic ovarian mass. *J gynecol oncol* 22: 244-252.
14. Ferraro S, Braga F, Lanzoni M, et al. (2013) Serum human epididymis protein 4 Vs carbohydrate antigen 125 for ovarian cancer diagnosis: a systematic review. *J Clin Pathol* 66: 273-281.
15. Dikmen ZG, Colak A, Dogan P, et al. (2015) Diagnostic performances of CA125, HE4, and ROMA index in ovarian cancer. *Eur J GynaecolOncol* 36: 457-462.
16. Moore RG, Jabre-Raughley M, Brown AK, et al. (2010) Comparison of a novel multiple marker assay Vs the risk of malignancy index for the prediction of epithelial ovarian cancer in patients with a pelvic mass. *Am J Obstet Gynecol* 203: 228e1-228e6.
17. Bonfrer JM, Korse CM, Verstraeten RA, et al. (1997) Clinical evaluation of the Byk LIA-mat CA125 II assay: discussion of a reference value. *Clin Chem* 43: 491-497.
18. Lycke M, Kristjansdottir B, Sundfeldt K (2018) A multicenter clinical trial validating the performance of HE4, CA125, risk of ovarian malignancy algorithm and risk of malignancy index. *Gynecol Oncol* 151: 159-165.

19. Chang X, Ye X, Dong L, et al. (2011) Human Epididymis Protein 4 (HE4) as a serum tumor biomarker in patients with ovarian carcinoma. *Int J Gynecol Cancer* 21: 852-858.
20. Oranratanaphan S, Wanishpongpan S, Termrungruanglert W, et al. (2018) Assessment of diagnostic values among CA-125, RMI, HE4, and ROMA for cancer prediction in women with nonfunctional ovarian cysts. *Obstetrics and gynecology international* 2018: 7821574.
21. Molina R, JM Escudero, JM Auge, et al. (2011) HE4 a novel tumour marker for ovarian cancer: comparison with CA 125 and ROMA algorithm in patients with gynaecological diseases. *Tumor Biol* 32: 1087-1095.
22. Chan KKL, Chen CA, Nam JH, et al. (2013) The use of HE4 in the prediction of ovarian cancer in Asian women with a pelvic mass. *Gynecol Oncol* 128: 239-244.
23. Roy LCDB (2019) Validity of the risk of malignancy index in malignant ovarian tumours. *International Journal of Scientific Research* 8.

DOI: 10.36959/468/487