



Research Article

DOI: 10.36959/973/447

Enzyme Activity of Rice Plant and Toxic in Acid Sulfate Soil, Mekong Delta, the Natural Poison Resistance of Rice Plant

Ba Le Huy^{1*}, Phong Nguyen Tan^{2*}, Hoan Nguyen Xuan³ and Phuong Nguyen Vu Hoang⁴

¹Chairman of Science and Education, Faculty of Biology and Environment, Ho Chi Minh City University of Industry and Trade (HUIT), Vietnam

²Dean, Faculty of Biology & Environment, Ho Chi Minh City University of Industry & Trade (HUIT), Vietnam

³President of Ho Chi Minh City University of Industry and Trade (HUIT), Vietnam

⁴Department of Biology and Environment, Ho Chi Minh City University of Industry and Trade (HUIT), Vietnam



Abstract

Aim: Determining the Toxic resistance of the rice plant, through the activity of three enzymes in rice: Catalase, Peroxydase and Phosphatase in the leaves of rice. In addition, consider the role of Phosphatase in the fluctuations and the ability to accumulate P_2O_5 in soil and in plants.

Methods: Combining experiments in the room, in greenhouses, greenhouses and fields, on acid sulfate soil, maintaining high toxicity but not killing rice plants. The experiments had four hybridization replicates; Full statistical processing.

Results: 1) High concentrations of toxins in the soil environment strongly inhibited the activity of Phosphatase enzymes in the roots, in the indicator leaves, and especially for rice, especially for plants that are less resistant to Al^{3+} toxins; 2) Enzyme Phosphatase in leaves has the effect of providing energy for the plants to recover from poisoning. This proves that the antagonism between enzymes and toxins is evident, especially with Al^{3+} toxins; 3) To confirm that the enzyme Peroxydase in leaves plays an important role, against the harmful effects of toxins; 4) Phosphatase enzyme in leaves has the effect on the providing energy for the plants to recover from poisoning.

Keywords

Toxic Inon in soil solution, Toxic-resistant of Enzyme in rice varieties, Toxicity in acid sulphate soil, Catalase Enzyme, Phosphatase Enzyme, Peroxidase Enzyme in rice: Acid sulfate soil, Mekong Delta

Abbreviation

TCVN: Việt Nam Standard; DBSCL: Mekong Delta: E/p: Enzyme Activity; R: Correlation coefficients; M: Characterization of Enzyme Activity in Rice

Introduction

Research of Le Huy Ba, et al. carry-out by [1-3] a series of experiments including: Step 1, Group testing of rice varieties including 15 varieties; In step 2, experiment comparing varieties to select alum tolerant varieties. Step 3, after identifying alum-tolerant varieties: Ca Dung variety (local) strong resistance to toxicity. Oppositely, IR38 variety is weakly to toxicity resistant. On other hand, they have found that, the toxic ions in acid sulphate soil environment (ex: Al^{3+} , Fe^{2+} , Fe^{3+} and SO_4^{2-}) there are coplicatly dynamics [1-4].

At this point, the new question that needs to be answered is the effect of toxins on the enzymes in plants and *vice versa*, the role of the enzyme has no role to resist poison (how effect

***Corresponding authors:** Ba Le Huy, Chairman of Science and Education, Faculty of Biology and Environment, Hochiminh City University of Industry and Trade (HUIT), Vietnam;

Phong Nguyen Tan, Dean, Faculty of Biology & Environment, Ho Chi Minh City University of Industry & Trade (HUIT), Vietnam

Accepted: September 23, 2024

Published online: September 25, 2024

Citation: Huy BL, Tan PN, Xuan HN, et al. (2024) Enzyme Activity of Rice Plant and Toxic in Acid Sulfate Soil, Mekong Delta, the Natural Poison Resistance of Rice Plant. J Rice Res Dev 6(1):466-483

on this activity of enzymes in rice leaves). That means, what role of this enzyme to help rice plant to resistant influence of the toxicity. So that, we carry out some experiences (*in vitro*, in greenhouse, and in the yield experiment together with the practical survey).

Materials and Research Methods

Material

Whole acid sulphate soil (not disturbed by cultivation): Heavy alkaline soil with pH = 3.6-4.1; Al^{3+} = 500-800 ppm, although the plants do not die but the toxicity is also very heavy, from which people choose resistant alum varieties [5-7].

Vietnamese rice varieties

- Select a selection of 15 new rice varieties including new varieties (IR38) and local varieties (Ca Dung)
- Compare and select toxicity-tolerant varieties: Including good alum-tolerant rice varieties, weak alum-resistant rice varieties (rice varieties living on alum soil during the research process).
- The indicator tree is also a control plant: *Elocharis dulcis*: Take normal growing plants, transplant into pots with alum soil.
- With the control variant of soil: Neutral alluvial soil (Tien river) is not alum or toxic.

So we have 2 control variant: With plants: Good alum-tolerant rice transplantation compared to poor alum-resistant rice and compared with indicator plants.

The toxicity and tolerance test of rice increased the toxicity levels (Fe^{2+} , Al^{3+} , SO_4^{2-} content increased gradually compared to the control (alluvial soil: non-toxic).

Experimental arrangement in pots placed in a greenhouse

- Arrangement in pots (diameter $r = 30$ cm, height: 50 cm, porcelain and porcelain materials), put in greenhouses (Poly Ethylene)
- Experiment with 5 treatments, 4 replications, sequentially according to latin cell, the level of toxicity: increasing, with treatments:
 - With Al^{3+}
1. $\text{Al}^{3+} = 0$ ppm, 2. $\text{Al}^{3+} = 135$ ppm, 3. $\text{Al}^{3+} = 500$, 4. $\text{Al}^{3+} = 800$ ppm and 5. $\text{Al}^{3+} = 1000$ ppm,
 - With Fe^{2+} :
1. $\text{Fe}^{2+} = 0$ ppm, 2. $\text{Fe}^{2+} = 600$ ppm; 3. $\text{Fe}^{2+} = 800$ ppm; 4. $\text{Fe}^{2+} = 900$ ppm; 5. $\text{Fe}^{2+} = 1000$ ppm.
- Maintaining sufficient alkaline water to flood the soil (5-7 cm)

Arranging field experiments:

- Rice plants: On Nhi Xuan acid sulphate soil, plot: 25 m², soil kept unchanged, no plowing, arranged Experiment 4 times, according to random sequential plot.

- Location: Whole alum soil belongs to Nhi Xuan Farm, Nhi Xuan commune, Binh Chanh district, Southwest, Ho Chi Minh City.
- Maintaining alum toxicity by adjusting the natural alum water in the area
- Adjust the toxicity through 2 periods: 1- high and 2- low to identify the reaction of rice physiology and biochemistry: The activity of 3 enzymes.

Tracking criteria:

- Growth and development of rice, especially 3-4 stages of true leaves (roots, stems, leaves...)
- Monitoring the development and toxicity of control acid sulphate soil eco-indicator plant (*Eleocharis dulcis*)
- Analysis of activity enzyme of rice plant: Take the second leaf (strongest activity leaf)

Analysis criteria:

- Soil and water environment: pH, Fe^{2+} , Al^{3+} , SO_4^{2-} , PO_4^{3-} by standard method (TCVN 2013)
- Activation of enzymes in functional leaves in rice: Amylase, Catalase, Peroxydase, Phosphatase Bio-analysis Method (Standar VN 2009)

Results

The effect of aluminum toxicity (Al^{3+}) and iron (Fe^{2+}) in soil on the growth of rice

Al^{3+} can be toxic even at low concentrations of 1-2 ppm (D. Dent- 1986, Le Huy Ba-1982) observed and analyzed on acid sulfate soil, said on the field when dry soil has a layer of $\text{Al}_2(\text{SO}_4)_3$ emerges in white, light and porous masses; When it is wet, it will contain mucus which contains 4.26% (Al^{3+}) and 38.4% (SO_4^{2-}). Al^{3+} has symptoms that are not always easy to recognize. For rice, the symptoms are expressed right at the roots (deformed, dented and brittle roots, although the color of the roots is not as black as iron poisoning), if the aluminum toxicity is high, the short rice roots will lose their hairs and die.

Le Huy Ba's research showed that at a concentration of Al^{3+} at 135 ppm beginning effect. While in the yield experiment, concentration of Al^{3+} at 500 ppm will effect on young rice plant (at rice plant have 3 real leaf). Thus, influence of toxic ion is dependent on environmental condition.

In addition, when Al^{3+} and Fe^{2+} exist simultaneously, the toxicity of soil also increases much more than normal [3].

In acid sulphate soils, Fe^{2+} , Fe^{3+} combine with S^{2-} to form FeS compound which attaches tightly to rice roots, impeding nutrient uptake in the soil as well as affecting respiration, causing rice roots to die.

The effect of Al^{3+} on young rice plants: Al^{3+} is a dangerous poison in the true tricuspid phase: At the concentration of $\text{Al}^{3+} = 135$ ppm in nutrient Konsac solution has begun to have an effect, especially for poorly tolerated varieties, and at 150

ppm there is a sign of death. And, up to 600 ppm die a lot. At 1,000 ppm in 20 days, the rice dies as softly as it is boiled water. It is worth noting that the roots are poisoned although the hairs are shed a lot, the roots are atrophied but the color is still white.

The influence of Fe^{2+} on young rice plants: At $\text{Fe}^{2+} = 150$ ppm, the effect has begun to affect and at 600 ppm, it is lethal to the less poisonous young rice varieties. However, manifestations in black roots and high mortality rates. Good poison tolerant varieties have a higher tolerance level, but if Fe^{2+} is at 1,000 ppm, the mortality will occur in both rice varieties very quickly. When the rice plant has Fe^{2+} poisoning, it affects the roots (the number of root hairs is reduced). Unlike Al poisoning, iron root poisoning is not white but becomes black, as a result of H_2S has an effect with Fe to form FeS, attaching to the outer layer of the rice roots, Fe^{2+} has decreased but still hinders domestic exchange processes. Moreover, the tips of the roots often become parrots in one piece, while the leaves have many spots similar to the disease of fire injection.

Compare the effects of Al^{3+} and Fe^{2+} toxins: At $\text{Fe}^{2+} = 600$ ppm, $\text{Al}^{3+} = 135$ ppm rice was damaged. On the other hand, when the concentration of $\text{Fe}^{2+} = 600$ ppm, $\text{Al}^{3+} = 135$ ppm, the harmful effects become many times greater than each

individual toxin. This is evident in all indicators and in turn goes from less poisoned to medium tolerant varieties to fairly toxic ones.

Effect of toxic concentration in acid sulphate environment on enzyme activity of Catalase enzyme and indicator

The results in Figure 1 indicate that the activity of this enzyme depends primarily on the growth period: They tend to decrease in both acid sulphate soils and in alum soil; Then came the influence of soil environmental toxins. On the other hand, we see that in acid sulfate environment, enzyme activity of Catalase is always lower than that on non-acid alluvial soil (average data of 8 varieties). This comment is at the confidence level with $t = 4,936$ and $t_{0.99} = 2,977$. This effect is more significant when we compare between alum-resistant varieties like Ca Dung and alum-resistant varieties like IR38, (Figure 1) also with 95% confidence.

While on alluvial soil, the enzyme activity of Enzyme Catalase in IR38 is always higher than that of Ca Dung, whereas in acid sulphate soil, the opposite is true: when toxic concentration increases (days 35 and 102), activity decreases (toxin). In soil, it is higher $\text{Al}^{3+} = 800\text{-}900$ ppm, $\text{SO}_4^{2-} = 0.3\text{-}0.35\%$, $\text{pH}_{\text{H}_2\text{O}} = 3.5\text{-}3.6$ (Figure 2).




Figure 1: Effects of toxic substances in the soil on the activity enzyme Catalase compares between non-toxic soils and acid sulphate soils.




Figure 2: Effect of toxic concentration in acid sulfate environment to enzyme activity of Catalase enzyme in indicator plant (*Eleocharis dulcis*) and rice plants.




Figure 3: Effect of toxins in soil environment on Peroxidase activity in rice roots and indicator plants.

Effect of toxic substances in acid sulphate soil environment on enzyme activity Peroxydase in indicator leaves (*Eleocharis dulcis*) and rice plants

Unlike Catalase, when the toxicity in acid sulphate environment increased (at 21, 60 days after sowing), the activity of the Peroxydase enzyme in rice leaves increased again (Figure 3). This rule is confirmed with high confidence $t = 3,400 > t_{0,99} = 2,977$.

The study results showed that the activity of the enzyme Peroxydase in the leaves depends on the growth period of the lua: High in the middle stage (60-70 days). On the other hand, it also depends on the characteristics of plants and varieties: The *Eleocharis dulcis* and Ca Dung alum tolerant varieties are always more active and more clearly expressed against the toxin than the less tolerant varieties, alum IR38.

Compared with the activity in non-alum alluvial soil, when the toxicity increased, Ca Dung variety had this activity increased by +280% (compared to the control), while the IR38 variety increased only +247% (compared to the control). This was data of *Eleocharis dulcis* - plant indicator for acid sulfate

Table 1: Toxic in acid sulphate soil environment.

Toxic in alum soil environment	high	low
Al ³⁺ (ppm)	805	465
Fe ²⁺ (ppm)	898	609
SO ₄ ²⁻ (%)	0.32	0.13

soil, supplemented and confirmed with absolute confidence (99%). That makes us appreciate, the right role of Peroxydase with alum poison resistance (Table 1).

In contrast, in the case of leaves, the peroxydase enzyme in plant roots living on acid sulfate soil was strongly inhibited of toxins in the soil environment (Figure 4). That is, when the toxicity in the soil increases, this enzyme activity in the roots of the leaves is less resistant to alum and alum tolerance, including in the grass roots. On the other hand, the enzyme activity in the root plays a small role: releasing energy, providing for lua, after being poisoned: For example, on the 87th day when the toxin decreased, this enzyme activity increased by +63% with Ca Dung and +40.1% with IR38.




Figure 4: Effects of toxins in soil environment on Peroxidase activity in rice roots (Ca Dung, IR38, *Eleocharis dulcis*).

Thus, according to the above result, the Enoxydasa enzyme in leaves plays a role in promoting Nang and Lua plants to fight against invading toxins. However, this role only appears to be within a certain limit of the level of toxicity in the soil environment and the characteristics of different varieties (Table 2).

Data in Table 2, show that the activity of this enzyme is significant in the limit of toxins in the soil environment:

- For Nang grass indicator plants that can tolerate toxic substances in soil at the highest level with:

$Al^{3+} \leq 950 \pm 31$ ppm, $Fe^{2+} \leq 907 \pm 44$ ppm, $SO_4^{2-} \leq 3209 \pm 48$ ppm.

- For Ca dung varieties, resistant to alum well, at the level:

$Al^{3+} \leq 890$ ppm, $Fe^{2+} \leq 842$ ppm, $SO_4^{2-} \leq 0.26\%$

- For poorly resistant alum like IR38, it is only at level:

$Al^{3+} \leq 750$ ppm, $Fe^{2+} \leq 676$ ppm, $SO_4^{2-} \leq 0.28\%$.

Correlation between toxins and P_2O_5 in the soil environment and enzyme activity Peroxydase in plants.

The toxic effect on this enzyme activity and the interaction between them is partly reflected through the correlation coefficient (R).

In the above correlation coefficients, it is worth noting:

the positive correlation between this Enzyme activity in the leaves of Nang grass and $R = +0.8432$. At the same time, we also found a positive correlation between the activity of the leaves of Ca Dung alum-tolerant alum with the concentration of Al^{3+} in the soil environment with $R = +0.802$. On the other hand, it should be noted the inverse correlation between this enzyme activity in alum-tolerant lua (IR38) with Al^{3+} content in soil environment with $R = -0.859$. It is the characteristics of alum-resistant plants different from the alum-resistant plants (Table 3).

Toxic effect in acid sulphate soil environment on enzyme Phosphatase activity in indicator and rice plants.

Influence Phosphatase in indicator leaves and rice plants: Toxicity in the acid sulphate environment not only limits the activity of enzyme Catalaza, Peroxydase in the roots, but also limits the activity of Phophatase enzyme in the leaves of the indicator plants and the lua. This is shown when the toxicity increased twice, this enzyme activity decreased at -21.2% and -53.6%, with Ca Dung variety; -40.7% and -69.6% for the IR38 variety, while for the Nang grass indicator plants, there is a decrease of -45% and -18%.

Equally interesting is the discovery of the role of Peroxydase in the leaves, where this Phophatase enzyme plays an important role in helping the plant recover after poisoning.

It is confirmed that this rule depends on the type and variety characteristics. They become significant in the case

of Ca Dung alum and indicator plants. For example, on day 87 when the toxicity in the soil decreased from low to high, this activity in the grass indicator increased by +59.9% and especially in alum-tolerant plants Ca dung increased by +643%, while the same was tolerated. Alum also increased but lower, at only +314% compared to the control. Another feature to note is that this activity also depends on the growth time of the later stages than the previous period.

Affects the Phophatase in rice roots: Unlike the leaves, the enzyme Phophatase in roots has very weak activity (Table 4). A comparison between the two varieties shows that this enzyme activity in the roots of Nang grass is a poison-tolerant adaptive wild plant that has a greater activity than alum-tolerant lua (Ca Dung). Alum tolerant lua root is more active than alum-resistant lua (IR38).

This is manifested in the inequality:
 $M \text{ (Eleocharis Dulcis)} > M \text{ (Ca Dung)} > M \text{ (IR38)}$
(M: Phosphatase enzyme activity in roots)

Correlation between toxins and P_2O_5 with Phosphatase: The data in Table 5 show that there is a negative and tight correlation between the concentration of Al^{3+} toxin in the soil and the Phophatase activity in the leaves, especially for the alum tolerant variety less than IR38. This proves that plants have mobilized Peroxidase into the body to support the vitality of trees. On the other hand, Phosphatase is inversely and tightly correlated with the amount of Al^{3+} in plants. This proves that there is an inverse and opposite reaction that inhibits each other between the toxin and Peroxidase

Table 2: Limiting the toxic effect of Enoxydase enzyme in plant leaves to toxic substances in the environment, on the 25th day after planting. (Enzyme activation unit: E/p)

Total concentration of toxic substances in the environment (ppm)				Plant type		
Variant	Al^{3+}	Fe^{2+}	SO_4^{2-}	<i>Eleocharis dulas</i>	Ca Dung	IR38
1	100 ± 5	135 ± 4	1000 ± 18	1.65 ± 0.07	1.26 ± 0.07	1.10 ± 0.08
2	450 ± 12	467 ± 18	1500 ± 47	2.47 ± 0.08	1.38 ± 0.04	1.15 ± 0.06
3	750 ± 16	676 ± 24	2108 ± 86	3.28 ± 0.10	1.62 ± 0.05	1.25 ± 0.05
4	890 ± 24	842 ± 32	2606 ± 58	3.30 ± 0.04	1.92 ± 0.10	0.62 ± 0.04
5	950 ± 31	907 ± 44	3209 ± 48	3.85 ± 0.14	1.64 ± 0.06	0.13 ± 0.00
6	1085 ± 525	1050 ± 58	3693 ± 92	1.67 ± 0.12	0.10 ± 0.0	0.00 ± 0.00
7	Non-toxic soil environment (control)			1.51 ± 0.04	1.45 ± 0.03	1.00 ± 0.05

($t_{0.95} = 2,447$)

Table 3: Correlation coefficient (R) between the concentration of P_2O_5 toxins in the soil environment and Enzyme Peroxidase (M) activity in the plants.

Correlate	M - SO_4^{2-}		M - Fe^{2+}		M - Al^{3+}		M - P_2O_5	
	Leaves	Roots	Leaves	Roots	Leaves	Root	Leaves	Root
<i>Eleocharis Dulcis</i>	+0.7204	-0.2015	+0.6344	-0.2104	+0.8432	-0.3215	-0.3467	+0.5241
Cà Dung	+0.6843	-0.2254	+0.6278	-0.2447	+0.8023	-0.3254	-0.3670	+0.4632
IR38	+0.3452	-0.4568	+0.4312	-0.5649	+0.6174	-0.8592	-0.2891	+0.3434

- Note: M is the activity of Enzyme Peroxidase
 $R_{0.95} = 0.754$

Table 4: The influence of toxic substances in acid sulphate soil environment on the enzyme activity Phophatase in the roots indicates acid sulphate-resistant and acid-resistant rice plants.

The day after sowing	Toxic in soil environment	Acid sulphate soil			Non toxic soil		
		<i>Eleocharis Dulcis</i>	Type of rice plants		<i>Eleocharis Dulcis</i>	Type of rice plants	
			Cà Dung	IR38		Cà Dung	IR38
21	High	1.29	1.29	0.96	0.14	3.50	2.21
36	Low	0.42	0.34	Small	0.42	0.30	0.41
66	High	Small	Small	Small	Streaks	Streaks	Streaks
87	Low	Small	Small	Small	Streaks	Streaks	Streaks

Table 5: Correlation (R) between toxic concentration in soil environment and activity (A) of Phosphatase enzyme in alum-resistant and alum-resistant plants.

Plants	A - SO ₄ ²⁻	A -Fe ²⁺	A - Al ³⁺	A - P ₂ O ₅
Cà Dung	-0.3648	-0.5601	-0.6462 ***	+0.6441***
IR38	-0.5001	-0.6239	-0.8972	+0.4427

Note: A: Phophatase enzyme activity in plants (R_{0.95} = 0.6320)
That allows us to deduce that phosphorus and phophatase play a large role in the mechanism of plant poison resistance to toxins in the soil environment.
(***: *Eleocharis dulcis*, A: phosphatase activity)

and phosphatase. Particularly noteworthy is the correlation between this enzyme activity and the accumulation of free phosphorus of Cà Dung alum tolerant variety (R = +0.6441), with 99% confidence.

Conclusion

Thus, based on the results of experiments of many years of research from 1979 up to now, it is possible to come to the conclusion:

- High concentrations of toxins in the soil environment strongly inhibited the activity of Phosphatase enzymes in the roots, in the indicator leaves, and especially for rice, especially for plants that are less resistant to alum.
- Enzyme Phosphatase in leaves has the effect of providing energy for the plants to recover from poisoning. The activity of this enzyme is inversely correlated with the content of Al³⁺ in soil and Al³⁺ in plants. This proves that the antagonism between enzymes and toxins is evident, especially with Al³⁺ toxins.
- Highly toxic in alum soil environment inhibits enzyme activity Peroxidase in rice roots. Under these conditions, peroxidase in the root plays a small role in helping the plant recover.
- We have sufficient basis to confirm that the enzyme Enoxydase in leaves plays an important role, against

the harmful effects of toxins. When the toxin in the soil, in plants increased activity of this enzyme also increased higher than the control.

When the toxicity increases, the Peroxidase in the leaves will also increase
+280% (compared to the control); while the alum-resistant plants did not increase (compared to the control).

When the toxicity decreased, Peroxydase in alum-resistant plants was also higher than alum-resistant: +643% for alum-resistant rice compared to the control. The poorly resistant alum variety only increased +314% (compared to the control).

Interest of Conflict

No interest of conflict.

References

1. Le Huy Ba (1980) Khanoc determined to determine the number of rice in a hurry, the toxic compounds on the Nhi Xuan Alum Strip. 4.
2. Le Huy Ba (1981) Problems of Acid sulphate soil in Mekong Delta, Vietnam. (Book in Vietnamese) Hochiming City General Publish House of, Vietnam.
3. Le Huy Ba (2003) Problems of Acid sulphate soil in Mekong Delta, Viet Nam. (Book in Vietnamese). The Publish House of National Hochiming City, Vietnam.
4. Le Huy Ba (1984) Evolution of Al³⁺, Fe²⁺, and SO₄²⁻ ions in soil, water, and plants during flood receding in Lang Bien acid sulfate soil. Journal of Agricultural Science and Technology, May 1984.
5. Le Huy Ba (1996) Soil Ecology, Ho Chi Minh City University Publish House 2000.
6. Pons LJ (1973) Outline of genesis, characteristics, classification and improvement of acid sulphate soils. In: Processing of the international symposium on acid sulphate soils. Wageningen, August 13-20. ILRI publication, 18, vol.
7. Akbar M, Ponnampereuma FN (1982) Saline soils South and Southeast Asia as potential rice lands. In: Rice research strategies for the future. IRRI, Los Banos, Philippines.

DOI: 10.36959/973/447

